



US005760069A

United States Patent [19]

Lukas-Laskey et al.

[11] Patent Number: 5,760,069

[45] Date of Patent: Jun. 2, 1998

[54] METHOD OF TREATMENT FOR
DECREASING MORTALITY RESULTING
FROM CONGESTIVE HEART FAILURE

[75] Inventors: Mary Ann Lukas-Laskey,
Philadelphia; Robert Ruffolo, Jr.,
Spring City; Neil Howard Shusterman,
Wynnewood, all of Pa.; Gisbert
Spöner, Laudenbach; Klaus Strein,
Hemsbach, both of Germany

[73] Assignee: Boehringer Mannheim
Pharmaceuticals
Corporation-SmithKline Beecham
Corporation Limited Partnership #1,
Rockville, Md.

[21] Appl. No.: 483,635

[22] Filed: Jun. 7, 1995

[51] Int. Cl.⁶ A61K 31/40; A61K 31/54;
A61K 31/34; A61K 31/585

[52] U.S. Cl. 514/411; 514/423; 514/223.2;
514/223.5; 514/471; 514/175

[58] Field of Search 514/411, 423,
514/223.2, 223.5, 471, 175

[56] References Cited

U.S. PATENT DOCUMENTS

4,888,179	12/1989	Appelgren et al.	424/480
5,308,862	5/1994	Ohlstein	514/411
5,312,828	5/1994	Finkelstein et al.	514/381

OTHER PUBLICATIONS

J. of Cardiovascular Pharm; Senior, et al., "Effects of Carvedilol on Ventricular Arrhythmias", 1992, vol. 19, (Suppl. 1): pp. S117-S121.

J. of Cardiovascular Pharm; DasGupta, et al., "The Effects of Intravenous Carvedilol, A New Multiple Action Vasodilatory β -Blocker, in Congestive Heart Failure", 1991, vol. 18, (Suppl. 1): pp. S12-S16.

British J. of Urology; Caine, et al., "The Use of Alpha-adrenergic Blockers in Benign Prostatic Obstruction", 1976, vol. 48, pp. 255-263.

British J. of Urology; H.N. Whitfield, et al., "The Effect of Adrenergic Blocking Drugs on Outflow Resistance", 1976, vol. 47, pp. 823-827.

American J. of Cardiology; DasGupta, et al., "Value of Carvedilol in Congestive Heart Failure Secondary to Coronary Artery Disease", 1990, vol. 66, pp. 1118-1123.

Z. Kardiol; A. Buchwald, et al., "Acute Hemodynamic Effects of the Beta-blocker Carvedilol in Heart Failure", 1990, vol. 79, No. 6, pp. 424-428.

JACC; DiLenarda, et al., "Acute Hemodynamic Effects of Carvedilol Versus Metoprolol In Idiopathic Dilated Cardiomyopathy", 1991, vol. 17, No. 2, Abstract 142A.

Frontiers in CHF; D. Tepper, "Multicenter Oral Carvedilol Heart Failure Assessment", 1996, vol. 2, No. 1, pp. 39-40.

J. of Cardiovascular Pharm; DasGupta, et al., 1990, vol. 19, (Suppl. 1): pp. 562-567.

J. of Hypertension, C. Rosendorff, "Beta-blocking agents with vasodilator activity", 1993, vol. 11, (Suppl. 4): pp. S37-S40.

Cardiology; J. Lessem, et al., "Development of a Multication Beta-blocker", 1993, vol. 82, (Suppl. 3): pp. 50-58.
Drug Safety; W.J. Louis, et al., "A Risk-Benefit Assessment of Carvedilol in the Treatment of Cardiovascular Disorders", 1994, vol. 11, No. 2, pp. 86-93.

Drugs; McTavish, et al., "Carvedilol—A Review of its Pharmacodynamic and Pharmacokinetic Properties, and Therapeutic Efficacy", 1993, vol. 45, No. 2, pp. 232-258.
Circulation; H. Krum, et al., "Effects of Carvedilol, a Vasodilator- β -Blocker, in Patients with Congestive Heart Failure Due to Ischemic Heart Disease", 1995, vol. 92, No. 2, pp. 212-218.

CBS-TV; CBS Evenings News, Transcript, Jan. 27, 1993, 6:30-7:00pm.

CNBC; Steals and Deals, Transcript, Jan. 29, 1993, 8:30pm.
Circulation, DasGupta, et al., 1989, vol. 80, No. 4, (Suppl. II): pp. 116-117.


Drugs of Today; Ruffolo, et al., "Carvedilol (Kredex): A Novel Multiple Action Cardiovascular Agent", 1991, vol. 27, No. 7, pp. 465-492.

Primary Examiner—William R. A. Jarvis

Attorney, Agent, or Firm—Mary E. McCarthy; Stephen Venetianer; Edward T. Lentz

[57] ABSTRACT

A method of treatment using a compound of Formula I:



wherein:

R₁ is hydrogen, lower alkanoyl of up to 6 carbon atoms or aroyl selected from benzoyl and naphthoyl;

R₂ is hydrogen, lower alkyl of up to 6 carbon atoms or arylalkyl selected from benzyl, phenylethyl and phenylpropyl;

R₃ is hydrogen or lower alkyl of up to 6 carbon atoms;

R₄ is hydrogen or lower alkyl of up to 6 carbon atoms, or when X is oxygen, R₄ together with R₅ can represent —CH₂—O—;

X is a valency bond, —CH₂, oxygen or sulfur;

Ar is selected from phenyl, naphthyl, indanyl and tetrahydronaphthyl;

R₅ and R₆ are individually selected from hydrogen, fluorine, chlorine, bromine, hydroxyl, lower alkyl of up to 6 carbon atoms, a —CONH₂— group, lower alkoxy of up to 6 carbon atoms, benzyloxy, lower alkylthio of up to 6 carbon atoms, lower alkylsulphonyl of up to 6 carbon atoms and lower alkylsulphonyl of up to 6 carbon atoms; or

R₅ and R₆ together represent methylenedioxy; or a pharmaceutically acceptable salt thereof, alone or in conjunction with one or more other therapeutic agents, said agents being selected from the group consisting of ACE inhibitors, diuretics, and digoxin for decreasing mortality resulting from congestive heart failure (CHF) in mammals, particularly humans.

7 Claims, No Drawings